

RECEIVED
CENTRAL FAX CENTER

SEP 28 2007

In The Claims:

Please amend the claims as follows:

Claims 1-8 (canceled)

9. (previously presented) A cream for the treatment of dermatitis characterized in that the volume ratios of respective ingredients are Lightyellow Sophora Root (*Sophora flavescens* Ait.), 2.7 to 3.3%; Turmeric (*Curcuma aromatica* Salisb.), 1.8 to 2.2%; Magnolia Bark (*Magnolia officinalis* Rehd. Et. Wils.), 1.8 to 2.2%; Moutan Bark (*Paeonia suffruticosa* Andr.), 1.8 to 2.2%; Isatis Leaf (*Isatis tinctoria* L.), 0.9 to 1.1%; Borneo Camphor Tree (*Dryobalanops aromatica* Gaertn. f.), 0.9 to 1.1%; Baikal Skullcap (*Scutellaria baikalensis* Georgi.), 1.8 to 2.2%; Amur Cork Tree (*Phellodendron amurense* Rupr.), 1.8 to 2.2%; Angelicae Dahuricae Root (*Angelica dahurica* Benth. et Hook.), 0.9 to 1.1%; Lemon (*Citrus Limonum*), 0 to 3%; Smartweed (*polygonum cuspidatum* Sieb. et Zucc), 0 to 1.1%; Licorice (*Glycyrrhiza uralensis* Fishch.), 0 to 0.55%; Cnidii Rhizoma (*Ligusticum chuanxiong* Hort), 0.45 to 1.1%; Japanese Angelica Root (*Angelica sinensis* (Olive) Diels.), 0 to 0.55%; salicylic acid, 0.45 to 0.55%; resorcinol, 0.45 to 0.55%; mutton oil, 2.7 to 3.3%; alcohol, 2.7 to 3.3%; and white soft paraffin, 63 to 78%.

10. (previously presented) A cream for the treatment of dermatitis characterized in that the volume ratios of respective ingredients are Lightyellow Sophora Root (*Sophora flavescens* Ait.), 3%; Turmeric (*Curcuma aromatica* Salisb.), 2%; Magnolia Bark (*Magnolia officinalis* Rehd. Et. Wils.), 2%; Moutan Bark (*Paeonia suffruticosa* Andr.), 2%; Isatis Leaf (*Isatis tinctoria* L.), 1%; Borneo Camphor Tree (*Dryobalanops aromatica* Gaertn. f.), 1%; Baikal Skullcap (*Scutellaria*

baikcalensis Georgi.), 2%; Amur Cork Tree (*Phellodendron amurense* Rupr.), 2%; Angelicae Dahuricae Root (*Angelica dahurica* Benth. Et Hook.), 1%; Lemon (*Citrus Limonum*), 3%; Smartweed (*polygonum cuspidatum* Sieb. Et Zucc), 1%; Licorice (*Glycyrrhiza uralensis* Fishch.), 0.5%; Cnidii Rhizoma (*Ligusticum chuanxiong* Hort), 0.5%; Japanese Angelica Root (*Angelica sinensis* (Olive) Diels.), 0.5%; salicylic acid, 0.5%; resorcinol, 0.5%; mutton oil, 3%; alcohol, 3%; and white soft paraffin, 70.5%.

11. (previously presented) A cream for the treatment of dermatitis characterized in that the volume ratios of respective ingredients are Lightyellow Sophora Root (*Sophora flavescens* Ait.), 2.7 to 3.3%; Turmeric (*Curcuma aromatica* Salisb.), 1.8 to 2.2%; Magnolia Bark (*Magnolia officinalis* Rehd. et. Wils.), 1.8 to 2.2%; Moutan Bark (*Paeonia suffruticosa* Andr.), 1.8 to 2.2%; Isatis Leaf (*Isatis tinctoria* L.), 0.9 to 1.1%; Borneo Camphor Tree (*Dryobalanops aromatica* Gaertn. f.), 0.9 to 1.1%; Baikal Skullcap (*Scutellaria baikcalensis* Georgi.), 1.8 to 2.2%; Amur Cork Tree (*Phellodendron amurense* Rupr.), 1.8 to 2.2%; Angelicae Dahuricae Root (*Angelica dahurica* Benth. et. Hook.), 0.9 to 1.1%; Lemon (*Citrus Limonum*), 0 to 3%; Smartweed (*polygonum cuspidatum* Sieb. et Zucc), 0 to 1.1%; Licorice (*Glycyrrhiza uralensis* Fishch.), 0 to 0.55%; dimethylsulfoxide, 4.5 to 5.5%; salicylic acid, 0.45 to 0.55%; resorcinol, 0.45 to 0.55%; mutton oil, 2.7 to 3.3%; alcohol, 2.7 to 3.3%; and white soft paraffin, 60 to 73%.

12. (previously presented) A cream for the treatment of dermatitis characterized in that the volume ratios of respective ingredients are Lightyellow Sophora Root (*Sophora flavescens* Ait.), 3%; Turmeric (*Curcuma aromatica* Salisb.), 2%; Magnolia Bark (*Magnolia officinalis* Rehd. et. Wils.), 2%; Moutan Bark (*Paeonia suffruticosa* Andr.), 2%; Isatis Leaf (*Isatis tinctoria* L.), 1%;

Borneo Camphor Tree (*Dryobalanops aromatica* Gaertn. f.), 1%; Baikal Skullcap (*Scutellaria baikcalensis* Georgi.), 2%; Amur Cork Tree (*Phellodendron amurense* Rupr.), 2%; Angelicae Dahuricae Root (*Angelica dahurica* Benth. et. Hook.), 1%; Lemon (*Citrus Limonum*), 3%; Smartweed (*polygonum cuspidatum* Sieb. et Zucc), 1%; Licorice (*Glycyrrhiza uralensis* Fishch.), 0.5%; dimethylsulfoxide, 5%; salicylic acid, 0.5%; resorcinol, 0.5%; mutton oil, 3%; alcohol, 3%; and white soft paraffin, 66.5%.

13. (withdrawn) The cream for the treatment of dermatitis of claim 9, wherein the Lightyellow Sophora Root (*Sophora flavescens* Ait.), the Turmeric (*Curcuma aromatica* Salisb.), the Magnolia Bark (*Magnolia officinalis* Rehd. Et. Wils.), the Moutan Bark (*Paeonia suffruticosa* Andr.), the Isatis Leaf (*Isatis tinctoria* L.) and the Borneo Camphor Tree (*Dryobalanops aromatica* Gaertn. f.) serve as main components, the Cnidii Rhizoma (*Ligusticum chuanxiong* Hort) and the Japanese Angelica Root (*Angelica sinensis* (Olive) Diels.) serve as accelerating agents for permeation, the salicylic acid and the resorcinol serve as accelerating agents for keratinization, and ratios of the main components: accelerating agents for skin permeation : accelerating agents for keratinization are 53 to 89% : 8 to 38% : 6 to 10%.

14. (withdrawn) The cream for the treatment of dermatitis of claim 10, wherein the Lightyellow Sophora Root (*Sophora flavescens* Ait.), the Turmeric (*Curcuma aromatica* Salisb.), the Magnolia Bark (*Magnolia officinalis* Rehd. Et. Wils.), the Moutan Bark (*Paeonia suffruticosa* Andr.), the Isatis Leaf (*Isatis tinctoria* L.) and the Borneo Camphor Tree (*Dryobalanops aromatica* Gaertn. f.) serve as main components, the Cnidii Rhizoma (*Ligusticum chuanxiong* Hort) and the Japanese Angelica Root (*Angelica sinensis* (Olive) Diels.) serve as

accelerating agents for permeation, the salicylic acid and the resorcinol serve as accelerating agents for keratinization, and ratios of the main components: accelerating agents for skin permeation : accelerating agents for keratinization are 53 to 89% : 8 to 38% : 6 to 10%.

15. (previously presented) The cream for the treatment of dermatitis of claim 11, wherein the Lightyellow Sophora Root (*Sophora flavescens* Ait.), the Turmeric (*Curcuma aromatica* Salisb.), the Magnolia Bark (*Magnolia officinalis* Rehd. Et. Wils.), the Moutan Bark (*Paeonia suffruticosa* Andr.), the Isatis Leaf (*Isatis tinctoria* L.) and the Borneo Camphor Tree (*Dryobalanops aromatica* Gaertn. f.) serve as main components, the dimethylsulfoxide serves as an accelerating agent for permeation, the salicylic acid and the resorcinol serve as accelerating agents for keratinization, and ratios of the main components: accelerating agent for skin permeation : accelerating agents for keratinization are 53 to 89% : 8 to 38% : 6 to 10%.


16. (withdrawn) The cream for the treatment of dermatitis of claim 12, wherein the Lightyellow Sophora Root (*Sophora flavescens* Ait.), the Turmeric (*Curcuma aromatica* Salisb.), the Magnolia Bark (*Magnolia officinalis* Rehd. Et. Wils.), the Moutan Bark (*Paeonia suffruticosa* Andr.), the Isatis Leaf (*Isatis tinctoria* L.) and the Borneo Camphor Tree (*Dryobalanops aromatica* Gaertn. f.) serve as main components, the dimethylsulfoxide serves as an accelerating agent for permeation, the salicylic acid and the resorcinol serve as accelerating agents for keratinization, and ratios of the main components: accelerating agent for skin permeation : accelerating agents for keratinization are 53 to 89% : 8 to 38% : 6 to 10%.

No new matter has been added to the application by the amendments made to the claims.

Dated: 9-28-2007

Correspondence Address:
4 Venture, Suite 250
Irvine, CA. 92618
Tel.: (949) 660-0761

Respectfully submitted,
J.C. PATENTS

By: 
Jiawei Huang
Registration No. 43,330